

Nils Postel<sup>1</sup>, Stephan Schneeweiss<sup>2</sup>, Christoph Wyen<sup>3</sup>, Farhad Schabaz<sup>4</sup>, Olaf Degen<sup>5</sup>, Gordon Weinberg<sup>6</sup>, Michael Sabranski<sup>7</sup>, Kevin Ummard-Berger<sup>8</sup>, Kathrin Maria Dymek<sup>9</sup>, Bernd Westermayer<sup>10</sup>, Jenny Scherzer<sup>9</sup>

<sup>1</sup>Prinzmed, Munich, Germany; <sup>2</sup>Praxis Hohenstaufenring, Cologne, Germany; <sup>3</sup>Praxis Ebertplatz, Cologne, Germany; <sup>4</sup>MVZ Karlsplatz, Munich, Germany; <sup>5</sup>Universitätsklinikum Hamburg-Eppendorf, Hamburg, Germany; <sup>6</sup>Infektiologisches Zentrum Steglitz, Berlin, Germany; <sup>7</sup>Infektionsmedizinisches Zentrum Hamburg (ICH), Hamburg, Germany; <sup>8</sup>UBN/Praxis, Berlin, Germany; <sup>9</sup>ViiV Healthcare, Munich, Germany; <sup>10</sup>GlaxoSmithKline, Munich, Germany

## Background

- The URBAN cohort study (initiated in 11/2018) provides prospective real-world data regarding the effectiveness, patient-reported outcomes (PROs) and tolerability of using Dolutegravir (DTG) plus Lamivudine (3TC) in people living with HIV either as two tablets (DTG+3TC) or – after availability in 7/2019 – as single tablet regimen (STR; DTG/3TC).
- Here we present the characteristics of the study population and month-6 (M6) outcomes.

## Methods

- URBAN is a prospective, non-interventional, 3-year German cohort study in ART-naïve and pre-treated patients receiving DTG plus 3TC in accordance with the label.
- Inclusion criteria for M6 full analysis set (FAS) were a documented M6 follow-up (visit window 4.5-9 months) or premature discontinuation.
- PROs were assessed using validated questionnaires (HIV Symptom Distress Module [HIV-SDM] and Treatment Satisfaction [HIV-TSQ]).

## Outcomes

- Month-6 (M6) viral suppression defined as HIV-RNA level [cp/mL] <50 or 50-200 with subsequent HIV-RNA <50 in the effectiveness set (missing=excluded)
- Persistence on study and/or DTG/3TC estimated using Kaplan-Meier analysis
- Adverse drug reactions (ADRs) coded by MedDRA (Medical Dictionary for Regulatory Activities) using system organ class (SOC) and preferred terms (PT)
- PRO measures at baseline and M3: mean/median total HIV-SDM and HIV-TSQ scores and changes from baseline

## Results

### Study population

- Overall, 367 patients were enrolled in the URBAN cohort.
- M6 FAS included 307 patients (92% pre-treated, 93% men, median age 48 years).
- Of pre-treated participants, 33% had a history of >3 ART regimens. Baseline HIV-RNA was <50 cp/mL in 96% of pre-treated patients; median HIV-RNA in ART-naïves was 37,100 copies/mL. Baseline characteristics are shown in Table 1.

### Reasons for use of DTG/3TC


- Primary reasons for use of DTG plus 3TC (in >15%) were 'preference of 2-drug regimen (2DR)' (29%) and 'side effects of previous ART' (23%) in pre-treated, and 'preference of 2DR' (48%) and 'easiness to take' (16%) in ART-naïves (Figure 1).

Table 1. Baseline characteristics

	Pre-treated	ART-naïve
Sex, male, n (%) [N]	262 (93) [282]	24 (96) [25]
Age, years, median (IQR) [N]	49 (39 – 55) [282]	35 (26 – 46) [25]
Age ≥50 years, n (%) [N]	132 (47) [282]	5 (20) [25]
Body weight, kg, median (IQR) [N]	79 (70 – 90) [219]	68 (65-83) [25]
BMI, kg/m <sup>2</sup> , median (IQR) [N]	25 (23 – 28) [217]	23 (21 – 25) [25]
Treatment start with fix-dose DTG/3TC, n (%) [N]	123 (44) [282]	5 (20) [25]
HIV-1 RNA, cp/mL, median (IQR) [N]	<50 [277]	37,100 (5,100-66,550) [25]
HIV-1 RNA >100,000 cp/mL, n (%)	1 (<1)	1 (4)
HIV-1 RNA <50 cp/mL, n (%)	267 (96)	0 (0)
CD4 T-cell count, cells/μL, median (IQR) [N]	738 (544 – 942) [277]	475 (386 – 664) [25]
History of AIDS (CDC C), n (%) [N]	36 (13) [282]	0 (0) [25]
Time since HIV diagnosis, years (median, IQR) [N]	10 (5 – 16) [279]	0 (0 – 0) [25]
Time on ART, years (median, IQR) [N]	7 (4 – 12) [253]	n.a.
Pretreatment, n (%) [N]	[280]	
INSTI-based	231 (83)	
NNRTI-based	23 (8)	n.a.
PI-based	20 (7)	
PI/INSTI-based	6 (2)	
Most common comorbidities (>10%), n (%) [N]	[282]	[25]
Hypertension	67 (24)	0 (0)
Depression	54 (19)	3 (12)
Chronic kidney disease	39 (14)	0 (0)
Insomnia	31 (11)	1 (4)
Lipid disorders	31 (11)	0 (0)

IQR: interquartile range; n.a.: not applicable


Figure 1. Primary reasons for use of DTG/3TC



### Persistence on study and/or DTG/3TC and discontinuation reasons

- Estimated persistence on study/DTG/3TC through M6 was 95% (Figure 2).
- 5% of patients (n=15/307) discontinued the study; reasons were adverse drug reactions (ADRs; 3.3%), virologic failure (investigator's discretion) (0.7%; n=2 with <200 copies/mL [without new resistance mutations, in one patient INI resistance test was not performed]), patient wish (0.3%), withdrawal of consent (0.3%) and doctor's decision (0.3%).

Figure 2. Persistence on study and/or DTG/3TC (Kaplan-Meier analysis)




### Safety

- Until M6, 22 ADRs (grades 1-2) were documented in 17 patients (6%).
- No serious ADR was reported.
- In 10 patients (3%), ADRs led to DTG/3TC discontinuation (n=13; several ADRs per patient possible; most common ADR was depression (n=3 [1%])).
- The following body weight changes through M6 were reported:
  - Median weight changes were +2.0 kg (IQR: -3.0 kg - +8.0 kg; n=7) in ART-naïve patients and +1.0 kg (-1.0 kg - +3.3 kg; n=104) in pre-treated patients.
  - No ADR or discontinuation due to weight gain were reported.

### Effectiveness

- Viral suppression rates are depicted in Figure 3.
- Of note, in 94% of pre-treated patients with ≥1 HIV-RNA follow-up, HIV-RNA was continuously <50 cp/mL.

Figure 3. Virologic outcomes at M6 (effectiveness set: N=269; n=38/307 with missing data)



\*including n=4 with 50-200 cp/mL & subsequent HIV-RNA level <50 cp/mL; \*\*n=1 with confirmed 50-200 cp/mL; \*n=1 with 360 cp/mL at M6 and 1,100,000 cp/mL at baseline; \*\*n=2 with a single HIV-RNA measurement of 128 and 89 cp/mL, respectively; \*\*ADRs (adverse drug reactions) (MedDRA SOC terms; multiple ADRs per patient possible) leading to discontinuation in >1% of the effectiveness set (N=269); psychiatric disorders (3%), skin and subcutaneous tissue disorders (1%).

### Patient reported outcomes

- In pre-treated patients completing PRO questionnaires at both time-points, mean changes in HIV-SDM and HIV-TSQ were -3.1 (p<0.001) and +2.5 (p<0.001), respectively (Table 2).

Table 2. Patient-reported outcomes in patients completing baseline and month-3 (M3) questionnaires

	Pre-treated patients	ART-naïve patients
<b>HIV Symptom Distress Module (HIV-SDM)<sup>^</sup>, N</b>	194 of 282	13 of 25
Baseline total score; mean/median (IQR)	14.5/12.0 (5.0 - 22.0)	12.2/9.0 (3.0 - 15.0)
M3 total score; mean/median (IQR)	11.4/8.0 (2.0 - 17.0)	7.6/3.0 (1.0 - 7.0)
Change from baseline mean/median (IQR) <sup>^^</sup>	-3.1/-3.0 (-8.0 - +1.0)	-4.6/-3.0 (-9.0 - ± 0.0)
p-value (Wilcoxon signed-rank test)	<0.001	0.068
<b>HIV Treatment Satisfaction (HIV-TSQ)<sup>*</sup>, N</b>	193	17
Baseline total score; mean/median (IQR)	53.4/56.0 (50.0 - 60.0)	n.a.
M3 total score; mean/median (IQR)	56.0/58.0 (53.0 - 60.0)	54.6/56.0 (54.0 - 58.0)
Change from baseline mean/median (IQR) <sup>**</sup>	+2.5/±0.0 (±0.0 - +4.0)	n.a.
p-value (Wilcoxon signed-rank test)	<0.001	n.a.

<sup>^</sup>HIV-SDM: 20 items, range of total score 0-80; <sup>^^</sup>negative changes indicate improvement; <sup>\*</sup>HIV-TSQ: range of total score 0-60; <sup>\*\*</sup>positive changes indicate improvement; IQR: interquartile range

## Conclusion

- DTG/3TC showed a high acceptance in ART-naïve and pre-treated patients with a persistence of 95% until month 6.
- Viral suppression rates were 90% in ART-naïve and 94% in pre-treated patients.
- Patients reported significant improvements in symptom distress and treatment satisfaction after the first months of treatment.

### Acknowledgments

Special thanks to the participating patients and investigators of the URBAN study centers: GP Isarpraxis/Munich; ICH/Hamburg; IZ Steglitz/Berlin; Klinikum Osnabrueck/Osnabrueck; MEDCENTER/Weimar; MVZ Karlsplatz/Munich; Onkologie Mannheim/Mannheim; Praxis City Ost/Berlin; Praxis Cordes/Berlin; Praxis Ebertplatz/Cologne; Praxis Hohenstaufenring/Cologne; Praxis Jessen/Berlin; Praxis Kaiserdamn/Berlin; Praxis Knechten/Aachen; Praxis UBN/Berlin; Praxis Wuensche/Berlin; PRINZMED/Munich; UNI Hamburg/Hamburg; WIR/Bochum. Support in medical writing was provided by MUC Research. The study is sponsored by ViiV Healthcare, Germany.