

Awareness and perception of accuracy of the Undetectable=Untransmittable (U=U) message in people living with HIV/AIDS (PLWHA), in Italy and correlation with the level of confidence in reference physicians

Cingolani A¹; Tavelli A²; Calvino GV³; Maggiolo F⁴; Girardi E⁵; Cozzi-Lepri A⁶; Perziano A⁷; Camposeragna A⁸; Gagliardini R⁹; Nozza S¹⁰; Antinori A⁹; d'Arminio Monforte, A¹¹

¹Fondazione Policlinico Agostino Gemelli IRCCS, Catholic University of the Sacred Heart, Infectious Diseases Unit, Rome, Italy ; ²Icona Foundation, Milan, Italy ; ³ANLAIDS Onlus, Rome, Italy ; ⁴ASST Papa Giovanni XXIII, Division of Infectious Diseases, Bergamo, Italy; ⁵National Institute for Infectious Diseases 'Lazzaro Spallanzani' IRCCS, Epidemiology Unit, Rome, Italy ; ⁶Institute for Global Health UCL, CREME, London, UK ; ⁷Associazione Arcobaleno AIDS, Torino, Italy ; ⁸Coordinamento Nazionale Comunità di Accoglienza (CNCA), Roma, Italy ; ⁹National Institute for Infectious Diseases 'Lazzaro Spallanzani' IRCCS, HIV/AIDS Department, Rome, Italy ; ¹⁰San Raffaele Scientific Institute IRCCS, Department of Infectious Diseases, Milan, Italy ; ¹¹ASST Santi Paolo e Carlo, University of Milan, Infectious Diseases Unit, Milan, Italy for ICONA Foundation Study Group

BACKGROUND

The efficacy of ART on HIV transmission has been widely demonstrated (1-7).

As a consequence, the Prevention Access Campaign has been set up to spread the slogan "Undetectable = Untransmittable" (friendly reported within online social media as #UequalsU) with the purpose of raising HIV prevention awareness and reducing HIV stigma.

At present, a considerable number of studies have demonstrated a low level of awareness of the message among those involved, never more than 15%, with higher percentages in the case of people living with HIV (8-10).

Purpose of this study was to evaluate the perceived accuracy of U=U message among PLWHA, HIV-negative people with sexual risky behaviors (PWSRB) and infectious diseases physicians, to guide subsequent efforts and implementation of HIV prevention strategies.

METHODS

An Italian nationwide web-survey among ICONA cohort centers, Community-based voluntary test&counselling centers (CBVTC) and fast-track cities websites has been conducted.

Three different anonymous questionnaires (for physicians, PLWHA and PWSRB) were set-up.

In this analysis the awareness of U=U ("have you ever heard of") and the perception of accuracy of U=U [Likert scale from 1=completely inaccurate (low) to 4=completely accurate (high) were analyzed.

Logistic regression models have been fitted to investigate factors associated with the binary outcomes (i)awareness of U=U (Y/N) and (ii)perceived high accuracy of U=U (Y/N).

RESULTS

1121 participants filled the questionnaires: 397 PLWHA, 90 physicians, 634 PWSRB. Participants' characteristics are shown in Table1. Awareness of U=U message has been reported in 74%, 46% and 92% of PLWHA, PWSRB and physicians.

Accuracy of U=U message has been reported as 'high' in 80% of PLWHA, 66% of PWSRB and 79% of physicians. Physicians perceived that 11% of PLWHA have a high perception of U=U; 34% of PLWHA reported a definitive positive messages received from physicians. Among PLWHA, factors associated with the awareness of U=U were level of education (university vs lower AOR=1.77, 95%CI 1.03-3.04), being MSM/bisexual (vs heterosexual AOR=3.16, 95%CI 1.03-3.04), being on cART for 5-10ys (vs <5ysAOR=2.71, 95%CI 1.32-5.55) and age (40-50ys vs <40y AOR=0.47, 95%CI 0.24-0.93). Factors associated with perception of accuracy of message in the three groups are reported in Figure1.

FUNDING


ICONA Foundation is supported by unrestricted grants from Gilead Sciences, Janssen, MSD, Thera technologies and ViiV Healthcare

Table 1 – Main Participants' characteristics; A) ID (Infectious Diseases) physicians; B) PLWHA (people living with HIV/AIDS) C) PWSRB (people with sexual risky behaviors)

A) ID physicians (N=90)		C) PWSRB (N=634)	
Age, years, n(%)		Age, years, n(%)	
<40 years	49 (54.4)	<40 years	461 (72.7)
40-50 years	18 (20.0)	40-50 years	110 (17.3)
>50 years	23 (25.6)	>50 years	63 (9.9)
Gender, Male, n(%)	37 (41.1)	Gender, Male, n(%)	431 (68.0)
Nationality, Italian, n(%)		Nationality, Italian, n(%)	610 (96.4)
Italian Geographical Zone, n(%)		Italian Geographical Zone, n(%)	
Northern	46 (51.1)	Northern	415 (65.7)
Central	38 (42.2)	Central	115 (18.2)
Southern/Islands	6 (6.7)	Southern/Islands	102 (16.1)
Education, University, n(%)	90 (100.0)	Education, University, n(%)	392 (61.8)
Management of PLWHA, Yes, n(%)	73 (81.1)	Management of PLWHA, Yes, n(%)	2 (1-10)
Yrs. management of PLWHA, n(%)		N. of sexual partners, median(IQR)	
<10 years	39 (53.4)	Sexual Orientation, n(%)	
10-20 years	15 (20.6)	Heterosexual	224 (35.3)
>20 years	19 (26.0)	Bisexual	42 (6.6)
Number of PLWHA in care, n(%)		Homosexual	368 (58.0)
<100	36 (49.3)	Stable Sexual Partner, n(%)	
100-400	15 (20.6)	Yes, HIV-pos	40 (6.3)
>400	22 (30.1)	Yes, HIV-neg	272 (42.9)
		Last HIV test, n(%)	
		<6 months	242 (38.2)
		6-12 months	95 (15.0)
		>12 months	176 (27.8)
		Never done	121 (19.1)

B) PLWHA (N=397)	
Age, years, n(%)	
<40 years	122 (30.79)
40-50 years	124 (31.2)
>50 years	151 (38.0)
Gender, Male, n(%)	324 (81.6)
Nationality, Italian, n(%)	375 (94.5)
Italian Geographical Zone, n(%)	
Northern	235 (59.2)
Central	117 (29.5)
Southern/Islands	45 (11.3)
Education, University, n(%)	157 (39.6)
Years with HIV infection	
<5 years	112 (28.2)
10-5 years	98 (24.7)
>10 years	187 (47.1)
Years of cART	
<5 years	125 (31.5)
10-5 years	113 (28.5)
>10 years	159 (40.0)
HIV-RNA undetectable, Yes, n(%)	372 (95.6)
N. of sexual partners, median(IQR)	2 (1-10)
Sexual Orientation, n(%)	
Heterosexual	131 (33.0)
Bisexual	41 (10.3)
Homosexual	225 (56.7)
Stable Sexual Partner, n(%)	
Yes, HIV-pos	53 (13.3)
Yes, HIV-neg	154 (38.8)

Figure 1. Factors associated with perceiving accuracy of U=U message as 'high' identified by multivariable logistic regressions analyses separately for each group



CONCLUSIONS

- A low concordance between awareness and perception of accuracy in PLWHA and physicians, suggesting still insufficient certainty was reported
- More efforts should be implemented to spread the U=U message among subgroups who might benefit from targeted educational campaigns
- Dissemination of the message among PWSRB is far from being efficaciously implemented and should represent a priority for increasing knowledge and decreasing HIV stigma
- Selection bias of web-surveys cannot be overlooked.

REFERENCES

1. Attia S, AIDS. 2009;23(11):1397-404. 2. Das M, PLoS One. 2010;5(6):e11068.3. Cohen MS, N Engl J Med. 2011;365(6):493-505. 4. Cohen MS, N Engl J Med. 2016;375(9):830-9. 5. Eshleman SH, J Acquir Immune Defic Syndr. 2017;74(1):112-6. 6. Rodger AJ, JAMA. 2016;316(2):171-81. 7. Bavinton BR, BMC Public Health. 2014;14(1):917. 8. Card KG, J Acquir Immune Defic Syndr. 2018; 77(1):8-16. 9. Carter A, J Int AIDS Soc. 2015;18(1):20039.10. Bavinton BR, Lancet HIV. 2018 Aug;5(8):e438-e447

ACKNOWLEDGMENTS

Icona Foundation Study Group: BOARD OF DIRECTORS: A d'Arminio Monforte (President), A Antinori (Vice-President), M Andreoni, A Castagna, F Castelli, R Cauda, G Di Perri, M Galli, R Iardino, G Ippolito, A Lazzarin, GC Marchetti, G Rezza, F von Schloesser, P Viale. **SCIENTIFIC SECRETARY:** A d'Arminio Monforte, A Antinori, A Castagna, F Ceccherini-Silberstein, A Cozzi-Lepri, E Girardi, A Gori, S Lo Caputo, F Maggiolo, C Mussini, M Puoti, CF Perno. **STEERING COMMITTEE:** A Antinori, F Bai, A Bandera, S Bonora, M Borderi, A Calcagno, MR Capobianchi, A Castagna, F Ceccherini-Silberstein, S Cicalini, A Cingolani, P Cinque, A Cozzi-Lepri, A d'Arminio Monforte, A Di Biagio, R Gagliardini, E Girardi, N Gianotti, A Gori, G Guaraldi, G Lapadula, M Lichtner, A Lai, S Lo Caputo, G Madeddu, F Maggiolo, G Marchetti, E Merlini, C Mussini, S Nozza, CF Perno, S Piconi, C Pinnetti, M Puoti, E Quiros Roldan, R Rossotti, S Rusconi, MM Santoro, A Saracino, L Sarmati, V Spagnuolo, V Svicher, L Taramasso. **STATISTICAL AND MONITORING TEAM:** A Cozzi-Lepri, I Fanti, L Galli, P Lorenzini, A Rodano', M Macchia, A Tavelli. **COMMUNITY ADVISORY BOARD:** A Bove, A Camposeragna, M Errico, M Manfredini, A Perziano, V Calvino. **BIOLOGICAL BANK INMI:** F Carletti, S Carrara, A Di Caro, S Graziano, F Petroni, G Prota, S Truffa. **PARTICIPATING PHYSICIANS AND CENTERS:** Italy A Giacometti, A Costantini, V Barocci (Ancona); G Angarano, L Monno, E Milano (Bari); F Maggiolo, C Suardi (Bergamo); P Viale, V Donati, G Verucchi (Bologna); E Quiros Roldan, C Minardi, (Brescia); F Franzetti, C Abeli (Busto Arsizio); L Chessa, F Pes (Cagliari); B Cacopardo, B Celesia (Catania); J Vecchiet, K Falasca (Chieti); A Pan, S Lorenzotti (Cremona); L Sighinolfi, D Segala (Ferrara); P Blanc, F Vichi (Firenze); M Bassetti, E Pontali, A Alessandrini, N Bobbio, G Mazzarello (Genova); M Lichtner, L Fondaco (Latina); S Piconi, C Molteni (Lecco); A Chiopera, P Milini (Macerata); G Nunnari, G Pellicano (Messina); A d'Arminio Monforte, M Galli, A Lazzarin, G Rizzardini, M Puoti, A Gori, A Castagna, ES Cannizzo, MC Moioli, R Piolini, D Bernacchia, A Poli, C Tincati (Milano); C Mussini, C Puzzolante (Modena); P Bonfanti, G Lapadula (Monza); V Sangiovanni, I Gentile, V Esposito, G Di Flumeri, G Di Filippo, V Rizzo (Napoli); AM Cattelani, S Marinello (Padova); A Cascio, C Colomba (Palermo); D Francisci, E Schiaroli (Perugia); G Parruti, F Sozio (Pescara); MA Di Pietro, A Vivarelli (Pistoia); C Lazzaretti, R Corsini (Reggio Emilia); M Andreoni, A Antinori, R Cauda, A Cristaudo, C Mastroianni, V Mazzotta, S Lamonica, M Capozzi, A Mondini, A Cingolani, M Rivano Capparuccia, G Iaiani, A Latini, G Onnelli, MM Plazzi, G De Girolamo, A Vergori (Roma); M Cecchetto, F Viviani (Rovigo); G Madeddu, A De Vito (Sassari); B Rossetti, F Montagnani (Siena); A Franco, R Fontana Del Vecchio (Siracusa); C Di Giulii (Terni); GC Orofino, G Di Perri, S Bonora, M Sciandra (Torino); C Tascini, A Londero (Udine); V Manfrin, G Battagin (Vicenza); G Starnini, A Ialungo (Viterbo).